

- Handbook to Birds of Australia (Gould), vol. ii., 1865, p. 479-480.
 Philos. Trans (Sharpe), vol. 168 (extra vol.), 1879, p. 130.
 Mem. Mus. Comp. Zool., vol. xiii., 1884. Water Birds of North America (Ridgeway), vol. ii., p. 418.
 Brit. Mus. Cat. Birds (Salvin), vol. xxv., 1896, p. 364-366.
 Nest and Eggs of Aust. Birds (Campbell), 1901, p. 875.
 Synopsis Avium; Nouveau Manuel Ornithologie (Dulois), vol. ii., 1903, p. 1030.
 "Ibis" (8), vol. vi., No. 24, 1906, p. 674.
 "Ibis" (8), vol. vi. No. 21, 1906, p. 149.
 Monograph of the Petrels (Godman), pt. 1, 1907, p. 59-67, pl. 17-18.
 Birds of Australia (Mathews), vol. ii., pt. 2, 1912, p. 31-47, pl. 71-72.
 List of Birds of Australia (Mathews), 1913, p. 32.
 Aust. Avian Record, vol. ii., No. 5, 1914, p. 86, loc. cit. No. 7, 1915, p. 124-5.
 Aust. Avian Record, vol. iv., No. 4-5, 1920, p. 108; loc. cit. No. 2-3, 1920, p. 67.
 Proc. Calif. Acad. Sci (Loomis). Review of the Albatrosses, Petrels and Diving Petrels, vol. ii., pt. ii., No. 12, p. 65-66, 183, 1918.
 Manual Birds of Australia (Mathews and Iredale), vol. 1, 1921, p. 19-21, pl. 4, fig. 6-7; pl. 7, fig. 6.
 In addition of the above, a complete search has been made through the "Journal Fur Ornithologie," "The Avicultural Magazine," "The Auk," "The Ibis," and "The Emu."

New Sub-species of *Acanthiza nana*, Vigors and Horsfield

Acanthiza nana flava, sub-sp. nov.

By H. L. WHITE, R.A.O.U., &c., Belltrees, Scone.

Upper surface bright dark citrine, becoming yellowish on upper tail coverts; crown medal bronze; wings and tail edged yellowish citrine on outer webs; dark subterminal bar on tail; small frontal spot buff; ear coverts fuscous with fine white shaft-streaks; throat clay colour (deep buff), with fine white shaft-streaks; breast olive yellow on sides, tinged buff in centre; abdomen flanks and undertail coverts lemon chrome; bill brown; legs dark brown; eyes brown; length, 89 mm.; bill, 8 mm.; wing, 51 mm.; tail, 38 mm.; tarsus, 16 mm.

Type, male (425) in "H. L. White Collection," Nat. Mus., Melb., taken at Ravenshoe, near Herberton, North Queensland, June, 1922. *Co-type*, female (426), from the same locality, has the buff on the throat more restricted; wing, 45 mm.

This, the most northerly and the brightest race of *A. nana* yet discovered, comes from a region elevated over 3000 feet above sea level, where another interesting member of the genus is found, *viz.*, *A. pusilla katherina*, De Vis.